

Effects of Standardized Ileal Digestible Threonine to Lysine Ratio on Growth Performance of PIC Line 337 × 1050 Pigs¹

Rafe Q. Royall, Mike D. Tokach, Jason C. Woodworth, Joel M. DeRouchey, Robert D. Goodband, Jordan T. Gebhardt,² Carine M. Vier,³ Matthew Spindler,³ Uislei Orlando,² Luis Zaragoza,³ Ning Lu,³ Wayne Cast,³ Danielle F. Wilson-Wells,³ Julia P. Holen,⁴ and Alyssa M. Betlach⁴

Summary

The objective of this research was to evaluate the impact of varying SID Thr:Lys ratios on growth performance, removals, and mortality rates of late-nursery, grower, and finishing PIC 337 × 1050 pigs. In each experiment, pens of pigs were blocked by BW and randomly assigned to 1 of 5 dietary treatments in a randomized complete block design with 19 to 27 pigs per pen and 8, 7, and 7 replications per treatment in Exp. 1, 2, and 3, respectively. In Exp. 1, 987 pigs (initially 26.0 ± 0.70 lb) were used from 26 to 54 lb. In Exp. 2, 875 pigs (initially 95.5 ± 1.17 lb) were used from 95 to 155 lb. In Exp. 3, 824 pigs (initially 224.4 ± 1.85 lb) were used from 224 to 297 lb. Pens were randomly assigned to 1 of 5 dietary treatments with increasing SID Thr:Lys ratios at 53, 58, 62, 65, and 68% in Exp. 1 and 2, and 56.5, 60, 64, 68, and 72.5% in Exp. 3. Diets were corn-soybean meal-based. Diets with the lowest and highest Thr:Lys ratios were blended to achieve the target SID Thr:Lys treatments in each experiment. Between experiments, all pens of pigs were placed on a common diet for 23 (Exp. 1 and 2) and 32 d (Exp. 2 and 3) to provide opportunity for compensatory growth prior to initiation of the next experiment. In Exp. 1 (26 to 54 lb), ADG and final BW increased linearly ($P \leq 0.006$) while ADFI, Thr intake/d, and Thr intake/kg of gain increased quadratically ($P \leq 0.001$). Overall, F/G improved (quadratic, $P \leq 0.001$) as Thr:Lys ratio increased. Additionally, Lys intake/d increased (quadratic, $P < 0.001$) while Lys intake/kg of gain decreased (quadratic, $P < 0.001$) with increasing Thr:Lys ratio. The quadratic polynomial (QP) model predicted greater than 68% SID Thr:Lys was required for ADG from 26 to 54 lb, while a QP model suggested that minimum F/G was achieved at 62.1% SID Thr:Lys. In Exp. 2 (95 to 155 lb), ADG, final BW, Thr intake/d, and Thr intake/kg of gain increased (linear, $P \leq 0.05$) and F/G improved (linear, $P = 0.030$) as dietary Thr:Lys increased. Moreover, Lys intake/kg of gain decreased (linear, $P = 0.023$)

¹ The authors appreciate Swine Vet Center (St. Peter, MN) and Genus PIC (Hendersonville, TN) for providing technical assistance for these studies.

² Department of Diagnostic Medicine/Pathobiology, College of Veterinary Medicine, Kansas State University.

³ Genus PIC, Hendersonville, TN.

⁴ Swine Vet Center, St. Peter, MN.

with increasing Thr:Lys ratio. For model analysis, QP models suggested optimum ADG and F/G were achieved at levels greater than 68% SID Thr:Lys. However, similar fitting broken-line quadratic (BLQ) and broken-line linear (BLL) models predicted no further improvement to F/G and ADG beyond 61 and 67% SID Thr:Lys, respectively. In Exp. 3 (224 to 297 lb), increasing SID Thr:Lys increased (linear, $P \leq 0.001$) Thr intake/d and Thr intake/kg of gain. In addition, increasing SID Thr:Lys ratios tended ($P \leq 0.086$) to quadratically increase ($P \leq 0.086$) ADFI and BW of pigs at the second marketing event. However, no other response criteria were impacted ($P \geq 0.10$) by dietary Thr:Lys. Due to a lack of ADG and F/G responses, prediction models were not developed. In summary, these results suggest the optimal SID Thr:Lys level for 26- to 54-lb pigs is 62.1% for feed efficiency and greater than 68% for ADG. From 95 to 155 lb, the requirement was predicted at or above 61 and 67% SID Thr:Lys for F/G and ADG, respectively. However, with the variation in response criteria in Exp. 3 (224 to 297 lb), we were unable to statistically define a requirement estimate.

Introduction

Threonine is an essential amino acid for pigs, which has been categorized as the first limiting amino acid for maintenance and development of intestinal tissue.⁵ An increase in the Thr requirement of pigs potentially occurs as pigs grow due to increased growth of intestinal tissue coupled with increased production of Thr-rich proteins (i.e., mucin).⁶ With improvements in modern swine genetics, it is critical to continuously re-evaluate the established nutrient requirements to optimize genetic potential for growth performance. Therefore, the objective of this study was to evaluate the impact of SID Thr:Lys ratios on growth performance, removals, and mortality rates of PIC 337 × 1050 late-nursery, grower, and finishing pigs.

Procedures

The protocol for this experiment was approved by the Kansas State University Institutional Animal Care and Use Committee. A commercial research barn located in south-central Minnesota was used to conduct the study. The barn had slatted concrete flooring, deep pits for manure storage, and was naturally ventilated. Pens contained a 3-hole stainless steel dry self-feeder (Thorpe Equipment, Thorpe, WI) and a 1-cup waterer to provide *ad libitum* access to feed and water. These experiments were conducted from May to September 2022.

Animals and diets

In Exp. 1, 987 pigs (PIC 337 × 1050; initially 26.0 ± 0.70 lb) were used from 26 to 54 lb. In Exp. 2, 875 pigs (initially 95.5 ± 1.17 lb) were used from 95 to 155 lb. In Exp. 3, 824 pigs (initially 224.4 ± 1.85 lb) were used from 224 to 297 lb. Pens were randomly assigned to 1 of 5 dietary treatments with increasing SID Thr:Lys ratios of 53, 58, 62, 65, or 68% in Exp. 1 and 2, and 56.5, 60, 64, 68, or 72.5% in Exp. 3. Diets were corn-soybean meal-based. Diets containing low and high Thr:Lys ratios were blended to achieve the intermediate SID Thr:Lys treatment levels in Exp. 1 and 2. In Exp. 3, diets

⁵ Tolosa A. F., M. D. Tokach, R. D. Goodband, J. C. Woodworth, J. M. DeRouchey, and J. T. Gebhardt. 2022. Evaluation of increasing digestible threonine to lysine in corn-soybean meal diets without and with distillers dried grains with solubles on growth performance of growing-finishing pigs. *Transl. Anim. Sci.* 6:1-6.

⁶ National Research Council. 2012. *Nutrient Requirements of Swine: Eleventh Revised Edition*. Washington, DC: The National Academies Press. <https://doi.org/10.17226/13298>.

containing 56.5, 64, and 72.5% SID Thr:Lys ratios were formulated. The 56.5 and 64% SID Thr:Lys diets were blended to create the 60% SID Thr:Lys treatment, while the 64 and 72.5% diets were blended to create the 68% SID Thr:Lys treatment (Table 1). Threonine was the first-limiting AA, with Lys formulated to be approximately 10% below PIC recommendations in each experiment, while all other AA ratios were maintained above PIC requirement estimates. Treatment diets were fed for 21, 28, and 36 d in Exp. 1, 2, and 3, respectively. Between experiments, all pens of pigs were placed on a common diet for 23 and 32 d, between Exp. 1 and 2, and Exp. 2 and 3, respectively. To determine ADG, ADFI, and F/G, pens of pigs were weighed, and feed disappearance was recorded throughout each experiment. On d 17 and 23 of Exp. 3, the 4 heaviest pigs from each pen were removed and marketed. The remaining pigs were marketed 13 d later at the conclusion of the study.

Statistical analysis

Data were analyzed as a randomized complete block design for a one-way ANOVA using the GLIMMIX procedure of SAS (v. 9.4, SAS Institute, Inc., Cary, NC). Pen was considered the experimental unit, initial body weight served as a blocking factor, and treatment served as the fixed effect in the statistical model. Contrast coefficients were adjusted to account for unequal spacing in treatments. Results were considered significant with $P \leq 0.05$ and marginally significant with $P \leq 0.10$. Dose response curves were evaluated using quadratic polynomial (QP), broken-line linear (BLL), and broken-line quadratic (BLQ) models. The best-fitting model was selected using the Bayesian Information Criterion (BIC) with improved model fits accepted when BIC decreased at least 2.0.

Results and Discussion

Experiment 1

In 26- to 54-lb pigs, ADG and final BW increased (linear, $P \leq 0.006$) with increasing Thr:Lys ratio. Meanwhile, ADFI, Thr intake/d, and Thr intake/kg of gain increased, and F/G improved (quadratic, $P \leq 0.001$) as Thr:Lys ratio increased. Additionally, Lys intake/d increased (quadratic, $P < 0.001$) while Lys intake/kg of gain decreased (quadratic, $P < 0.001$) with increasing Thr:Lys ratio. There was no difference between dietary treatments on the percentage of removals, mortalities, or total removals ($P > 0.10$).

A QP model predicted the requirement at levels greater than 68% SID Thr:Lys for ADG, while a QP model suggested that 62.1% SID Thr:Lys was required to achieve minimum F/G (Figures 1 and 2).

Experiment 2

In 95- to 155-lb pigs, ADG, final BW, Thr intake/d and Thr intake/kg of gain increased (linear, $P \leq 0.036$) as dietary Thr:Lys increased, while F/G improved (linear, $P = 0.030$). Moreover, Lys intake/kg of gain decreased (linear, $P = 0.023$) with increasing Thr:Lys. There was no difference between treatments on the percentage of removals, mortalities, or total removals ($P > 0.10$).

Quadratic polynomial models suggested optimum ADG and F/G were achieved at dietary levels greater than 68% SID Thr:Lys. Similar fitting BLQ and BLL models

predicted no further improvement to F/G and ADG beyond 61 and 67% SID Thr:Lys, respectively (Figures 3 and 4).

Experiment 3

In 224- to 297-lb pigs, increasing SID Thr:Lys ratio increased (linear, $P \leq 0.001$) Thr intake/d and Thr intake/kg of gain. In addition, increasing SID Thr:Lys ratios marginally increased (quadratic, $P \leq 0.086$) ADFI and BW of pigs at the second marketing event. However, no other response criteria were significantly impacted by dietary Thr:Lys ($P \geq 0.10$). Additionally, there was no difference across Thr:Lys treatments on the percentage of removals, mortalities, or total removals ($P > 0.10$). Models were not analyzed for Exp. 3 due to lack of observed significant differences for ADG or F/G.

In summary, these results suggest the optimal SID Thr:Lys level for 26- to 54-lb pigs is 62.1% for feed efficiency and greater than 68% for ADG. For pigs from 95 to 155 lb, the requirement was predicted to be at or above 65 and 67% SID Thr:Lys for F/G and ADG, respectively. However, statistical differences across dietary SID Thr:Lys levels for 224- to 297-lb pigs were not detected, thus models to establish requirements for growth performance criteria were not conducted.

Brand names appearing in this publication are for product identification purposes only. No endorsement is intended, nor is criticism implied of similar products not mentioned. Persons using such products assume responsibility for their use in accordance with current label directions of the manufacturer.

Table 1. Diet composition (as-fed basis)^{1,2}

Item	Exp. 1		Exp. 2		Exp. 3		
	Low	High	Low	High	Low	Medium	High
Ingredient, %							
Corn	64.86	64.67	78.39	78.17	84.72	84.66	84.64
Soybean meal	30.00	30.00	18.70	18.8	13.25	13.25	13.20
Choice white grease	1.45	1.45	---	---	---	---	---
Monocalcium P, 21% P	0.87	0.87	0.45	0.45	---	---	---
Limestone	1.00	1.02	1.18	1.18	1.00	1.00	1.00
Sodium chloride	0.61	0.61	0.55	0.55	0.56	0.56	0.56
L-Lys-HCl	0.36	0.36	0.28	0.28	0.21	0.21	0.21
DL-Met	0.23	0.23	0.11	0.11	0.05	0.05	0.05
L-Thr	0.03	0.20	---	0.12	---	---	---
Thr biomass ³	---	---	---	---	---	0.06	0.13
L-Trp	0.02	0.02	0.01	0.01	0.01	0.01	0.01
L-Val	0.13	0.13	0.04	0.04	0.01	0.01	0.01
Vitamin premix with phytase ⁴	0.25	0.25	0.15	0.15	0.10	0.10	0.10
Trace mineral premix	0.15	0.15	0.15	0.15	0.10	0.10	0.10
Copper sulfate	0.06	0.06	---	---	---	---	---
Total	100	100	100	100	100	100	100
Calculated analysis ¹							
SID AA, %							
Lys, %	1.15	1.15	0.81	0.81	0.63	0.63	0.63
Ile:Lys	56	56	59	59	63	63	63
Leu:Lys	119	119	131	131	149	149	149
Met and Cys:Lys	59	59	59	59	59	59	59
Thr:Lys	53	68	53	68	56.5	64.0	72.5
Trp:Lys	20	20	19.2	19.2	19.0	19.0	19.0
Val:Lys	70	70	70	70	73	73	73
His:Lys	36	36	37	37	40	40	40
NE, kcal/lb	1,111	1,111	1,121	1,121	1,142	1,142	1,142
SID Lys:NE, g/Mcal	4.70	4.70	3.28	3.28	2.50	2.50	2.50
CP, % ⁵	18.7	18.8	14.2	14.3	12.0	12.0	12.1
Ca, %	0.69	0.70	0.64	0.64	0.47	0.47	0.47
STTD P, %	0.44	0.44	0.33	0.33	0.26	0.26	0.26

¹Calculated analysis is based off nutrient profiles for ingredients listed in the NRC. (National Research Council. 2012. Nutrient Requirements of Swine: Eleventh Revised Edition. Washington, DC: The National Academies Press. <https://doi.org/10.17226/13298>.)

²Low and high Thr:Lys diets were blended to create treatment diets at 58, 62, and 65% SID Thr:Lys in Exp. 1 and 2. In Exp. 3, the low and medium diets were blended to create the 60% SID Thr:Lys treatment, and the medium and high diets were blended to create the 68% SID Thr:Lys treatment.

³THR Pro; CJ America-Bio, Downers Grove, IL.

⁴Axtra PHY (Dupont, Wilmington, DE) provided 325, 195, and 236 phytase units (FTU/lb) for an estimated release of 0.13, 0.12, and 0.14% STTD P, in Exp. 1, 2, and 3, respectively.

⁵CP = crude protein.

Table 2. Effects of increasing SID Thr:Lys on growth performance of 26- to 54-lb PIC 337 × 1050 pigs, Exp. 1^{1,2}

Item	SID Thr:Lys, %					SEM	P =	
	53	58	62	65	68		Linear	Quadratic
BW, lb								
d 0	26.4	25.8	26.0	25.6	26.3	0.70	0.532	0.109
d 21	51.0	52.2	53.1	52.9	53.9	0.90	0.006	0.775
d 0 to 21								
ADG, lb	1.16 ^b	1.21 ^{ab}	1.24 ^{ab}	1.25 ^{ab}	1.29 ^a	0.024	0.003	0.951
ADFI, lb	1.87 ^{ab}	1.72 ^c	1.72 ^c	1.74 ^{bc}	1.92 ^a	0.047	0.599	< 0.001
F/G	1.61 ^a	1.43 ^b	1.40 ^b	1.38 ^b	1.49 ^{ab}	0.046	0.002	< 0.001
Thr intake, g/d	5.34 ^c	5.39 ^c	5.76 ^{bc}	6.10 ^b	7.05 ^a	0.157	< 0.001	< 0.001
Thr intake, g/kg gain	10.18 ^b	9.86 ^b	10.30 ^b	10.70 ^b	12.05 ^a	0.333	< 0.001	0.001
Lys intake, g/d	10.09 ^{ab}	9.29 ^c	9.29 ^c	9.35 ^{bc}	10.36 ^a	0.250	0.644	< 0.001
Lys intake, g/kg gain	19.20 ^a	17.00 ^b	16.61 ^b	16.48 ^b	17.74 ^{ab}	0.544	0.002	< 0.001
Removals and mortality								
Removals, %	1.51	2.51	2.53	1.53	3.59	1.332	0.430	0.990
Mortality, %	0.28	0.28	0.29	0.00	0.00	0.437	0.997	0.998
Total, %	2.01	3.02	3.03	1.53	3.59	1.332	0.720	0.969

^{ab}Means within row with different superscripts differ ($P < 0.05$).

¹A total of 987 pigs (initial BW = 26.0 ± 0.70 lb) were used in a 21-d growth performance study with 22 to 27 pigs per pen and 8 replicates per treatment.

²ADG = average daily gain. ADFI = average daily feed intake. F/G = feed-to-gain ratio.

Table 3. Effects of increasing SID Thr:Lys on growth performance of 95- to 155-lb PIC 337 × 1050 pigs, Exp. 2^{1,2}

Item	SID Thr:Lys, %					SEM	P =	
	53	58	62	65	68		Linear	Quadratic
BW, lb								
d 0	95.4	95.4	95.5	95.5	95.5	1.17	0.841	0.958
d 28	151.1	151.3	152.0	153.9	154.3	1.76	0.036	0.507
d 0 to 28								
ADG, lb	1.98	2.00	2.01	2.09	2.08	0.037	0.028	0.676
ADFI, lb	4.61	4.52	4.63	4.64	4.69	0.068	0.212	0.321
F/G	2.32 ^a	2.27 ^{ab}	2.30 ^{ab}	2.22 ^b	2.26 ^{ab}	0.025	0.030	0.584
Thr intake, g/d	9.51	10.23	11.20	11.76	12.43	0.163	< 0.001	0.324
Thr intake, g/kg gain	10.59	11.30	12.29	12.41	13.21	0.130	< 0.001	0.711
Lys intake, g/d	17.96	17.66	18.04	18.10	18.29	0.266	0.210	0.337
Lys intake, g/kg gain	19.97	19.50	19.80	19.09	19.41	0.214	0.023	0.613
Removals and mortality								
Removals, %	0.00	0.00	0.00	0.00	0.58	0.576	0.992	0.993
Mortality, %	0.53	1.09	0.53	0.54	1.06	0.860	0.816	0.934
Total, %	0.59	1.19	0.59	0.60	1.73	0.993	0.614	0.730

^{ab}Means within row with different superscripts differ ($P < 0.05$).

¹A total of 875 pigs (initial BW = 95.5 ± 1.17 lb) were used in a 28-d growth performance study with 21 to 27 pigs per pen and 7 replicates per treatment.

²ADG = average daily gain. ADFI = average daily feed intake. F/G = feed-to-gain ratio.

Table 4. Effects of increasing SID Thr:Lys ratio growth performance of 224- to 297-lb PIC 337 × 1050 pigs, Exp. 3^{1,2}

Item:	SID Thr:Lys, %					SEM	P =	
	56.5	60	64	68	72.5		Linear	Quadratic
BW, lb								
d 0	224.7	224.1	223.7	224.4	224.4	1.85	0.916	0.434
d 36 ³	294.4	294.8	296.2	296.2	294.8	3.26	0.788	0.543
Marketing								
First marketing (d 17) ⁴	283.5	277.5	281.0	281.6	280.2	3.29	0.819	0.620
Second marketing (d 23) ⁴	293.3	294.6	297.1	297.3	287.7	3.71	0.434	0.086
Average ⁵	292.3	291.8	293.4	293.9	291.1	2.22	0.926	0.246
d 0 to 36								
ADG, lb	2.21	2.20	2.26	2.29	2.18	0.044	0.935	0.144
ADFI, lb	6.47	6.56	6.57	6.65	6.40	0.099	0.778	0.083
F/G	2.93	2.99	2.91	2.91	2.94	0.036	0.666	0.750
Thr intake, g/d	10.43 ^d	11.57 ^c	12.39 ^b	13.13 ^a	13.23 ^a	0.183	< 0.001	< 0.001
Thr intake, g/kg gain	10.39 ^d	11.63 ^c	12.07 ^c	12.69 ^b	13.40 ^a	0.147	< 0.001	0.039
Lys intake, g/d	19.69	19.91	19.96	20.20	19.46	0.304	0.789	0.096
Lys intake, g/kg gain	19.60	20.04	19.46	19.53	19.70	0.238	0.654	0.772
Removals and mortality								
Removals, %	0.00	0.00	0.80	0.00	0.38	0.974	0.990	1.000
Mortality, %	0.00	0.43	0.90	0.88	0.43	0.844	0.982	0.981
Total, %	0.00	0.51	2.17	1.07	1.04	1.329	0.985	0.985

^{ab}Means within row with different superscripts differ ($P < 0.05$).

¹A total of 824 pigs (initial BW = 224.4 ± 1.85 lb) were used in a 36-d growth performance study with 19 to 26 pigs per pen and 7 replicates per treatment.

²ADG = average daily gain. ADFI = average daily feed intake. F/G = feed-to-gain ratio.

³Final marketing event occurred on d 36.

⁴The 4 heaviest pigs were marketed from each pen at each marketing event.

⁵Weighted average final BW for all marketing events (d 17, 23, and 36).




Figure 1. Estimation of SID Thr:Lys requirements to maximize ADG for 26- to 54-lb PIC 337 × 1050 pigs, Exp. 1.

A total of 987 pigs (PIC 337 × 1050; initially 26.0 ± 0.70 lb) were used in a 21-d trial. Quadratic polynomial (QP), broken-line linear (BLL), and broken-line quadratic (BLQ) models were fit to estimate SID Thr:Lys level required to maximize ADG. The QP model resulted in the best fit, based on Bayesian Information Criterion (BIC), with a lower number being indicative of a better fit. The QP model predicted 95 and 100% of maximum ADG at greater than 68% SID Thr:Lys. The developed QP model equation for ADG was: $ADG = -0.00003 \times (SID\ Thr:Lys, \%)^2 + 0.01167 \times (SID\ Thr:Lys, \%) + 0.6244$.




Figure 2. Estimation of SID Thr:Lys requirements to minimize F/G for 26- to 54-lb PIC 337 × 1050 pigs, Exp. 1.

A total of 987 pigs (PIC 337 × 1050; initially 26.0 ± 0.70 lb) were used in a 21-d trial. Quadratic polynomial (QP), broken-line linear (BLL), and broken-line quadratic (BLQ) models were fit to estimate SID Thr:Lys level required to minimize F/G. The QP model resulted in the best fit, based on Bayesian Information Criterion (BIC), with a lower number being indicative of a better fit. The QP model predicted 95 and 100% of minimum F/G at 57.1 and 62.1% SID Thr:Lys. The developed QP model equation for F/G was: $F/G = 0.002845 \times (\text{SID Thr:Lys, \%})^2 - 0.3531 \times (\text{SID Thr:Lys, \%}) + 12.336$.




Figure 3. Estimation of SID Thr:Lys requirements to maximize ADG for 95- to 155-lb PIC 337 × 1050 pigs, Exp. 2.

A total of 875 pigs (PIC 337 × 1050; initially 95.5 ± 1.17 lb) were used in a 28-d trial. Quadratic polynomial (QP), broken-line linear (BLL), and broken-line quadratic (BLQ) models were fit to estimate SID Thr:Lys level to maximize ADG. The QP and BLL models resulted in the best fit, based on Bayesian Information Criterion (BIC), with a lower number being indicative of a better fit (BIC = -60.0 vs. -58.5, QP vs. BLL). The QP model predicted 95 and 100% of maximum ADG at 66.2 and greater than 68% SID Thr:Lys, respectively. The developed QP model equation for ADG was: $ADG = 0.000288 \times (SID\ Thr:Lys, \%)^2 - 0.0275 \times (SID\ Thr:Lys, \%) + 2.6277$. The BLL model predicted no further improvement beyond 67% SID Thr:Lys.




Figure 4. Estimation of SID Thr:Lys requirements to minimize F/G for 95- to 155-lb PIC 337 × 1050 pigs, Exp. 2.

A total of 875 pigs (PIC 337 × 1050; initially 95.5 ± 1.17 lb) were used in a 28-d trial. Quadratic polynomial (QP), broken-line linear (BLL), and broken-line quadratic (BLQ) models were fit to estimate SID Thr:Lys level to minimize F/G. The QP and BLQ models resulted in the best fit, based on Bayesian Information Criterion (BIC), with a lower number being indicative of a better fit (BIC = -81.3 vs. -79.5, QP vs. BLQ). The QP model predicted 95 and 100% of minimum F/G at levels greater than 68% SID Thr:Lys. The developed QP model equation for F/G was: $F/G = 0.000263 \times (\text{SID Thr:Lys, \%})^2 - 0.03654 \times (\text{SID Thr:Lys, \%}) + 3.5203$. The BLQ model suggested no further improvement in F/G beyond 61% SID Thr:Lys.