

TURFGRASS RESEARCH 2016



JULY 2016

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service

K-State Research and Extension is an equal
opportunity provider and employer.

Nitrous Oxide Emissions and Carbon Sequestration in Turfgrass: Effects of Irrigation and Nitrogen Fertilization (Year 1)

Ross C. Braun, Dale J. Bremer, and Jack D. Fry

Summary. In this study, annual nitrous oxide (N_2O) emissions were greatest in urea and least in untreated (no N) among treatments. Differences were negligible due to irrigation treatment. Irrigation levels may be decreased further in the final year to induce slight stress on the low irrigation treatment. All fertilizer treatments maintained acceptable quality, however the controlled-release fertilizer resulted in more consistent visual quality ratings compared to urea and untreated. Urea fertilizer had higher peak fluxes after fertilization and overall annual emissions than polymer-coated N-fertilizer. Thus, controlled released N fertilizers, such as polymer-coated urea, in turfgrass systems could potentially help mitigate N_2O emissions.

Rationale. Nitrous oxide (N_2O) is an important greenhouse gas implicated in global climate change. Turfgrass is typically fertilized with nitrogen (N) and irrigated while having the potential to emit N_2O at similar rates as other agricultural soils and thus, play an important role in atmospheric N_2O budgets. The development of management practices, such as slow-release N fertilizer and/or deficit irrigation, may mitigate N_2O emissions.

Objectives. To quantify the magnitude and patterns of N_2O emissions in turfgrass and determine how irrigation and N fertilization may be managed to reduce N_2O emissions.

Study Description. The study is being conducted under an automated rainout shelter near Manhattan, KS (Figure 1). By shielding rainfall from turfgrass, researchers can control the amount of water applied to plots. 'Meyer' zoysiagrass was sodded

View all turfgrass research reports online at: <http://newprairiepress.org/kaesrr>



June 4, 2013, and maintained at a 1-inch mowing height. During the summer (June - August) of 2014 and 2015, two irrigation treatments were applied including medium (72% reference evapotranspiration [ET_0] replacement) and medium-low (54% ET_0 replacement) (Figure 2). Three N-fertilization treatments included urea and polymer-coated N, both at 2 lb/1000 ft², and a control with no N applied. Because little drought stress was observed in the 54% ET_0 treatment in 2014, irrigation amounts in both treatments were reduced, specifically from 72 to 68% ET_0 replacement in the medium and from 54 to 33% ET_0 replacement in the medium-low treatment in 2015.

Nitrous oxide emissions were measured with static chambers placed over the turfgrass surface and then analyzed with gas chromatography (Figure 3). Measurements began on October 29, 2014 (DOY 302) and continued weekly-to-monthly until October 5, 2015 (DOY 278), concluding the first year of data. Irrigation and fertilization treatments will be repeated in 2016 for a second year of measurements through October 30, 2016. Carbon sequestration in the upper soil profile (0 to 12 inches) will be measured by sampling soil C at the end of the 3-year study; initial soil C was measured on August 28 2013. Ancillary measurements included soil moisture, temperature, nitrate, ammonium, visual quality, mowing frequency, and percent green cover using digital image analysis.

Results. Cumulative annual emissions of N_2O were significantly greatest in urea and least in untreated (no N) zoysiagrass among treatments. Annual emissions were 1.82, 2.09, 2.77 kg N_2O -N ha⁻¹ yr⁻¹ for untreated, polymer-coated urea, and urea, respectively (Figure 4). Annual emissions were similar to those reported in other turfgrass studies, which ranged from 1 to 3.85 kg N_2O -N ha⁻¹ yr⁻¹ across various turfgrass species and under different fertilization regimes (Bremer, 2006; Kaye et al., 2004; Lewis and Bremer, 2013). The percentage of applied N fertilizer emitted as N_2O in this study was 2.2% from poly and 2.9% from urea fertilizer. The highest fluxes and majority of emissions occurred in the summer because of the fertilization events and, presumably, higher soil temperatures. There were spikes after applications of urea fertilizer, but increases were much smaller with application of controlled release (poly) N fertilizer. Both urea and controlled release N fertilizer treatments resulted in higher turfgrass quality than the control; however, all three treatments maintained acceptable turfgrass quality during deficit irrigation treatments (Figure 5).

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service





References

- Bremer, D.J. 2006. Nitrous oxide fluxes in turfgrass: Effects of nitrogen fertilization rates and types. *J. Environ. Qual.* 1678-1685.
- Kaye, J.P., I.C. Burke, A.R. Mosier, and J.P. Guerschman. 2004. Methane and nitrous oxide fluxes from urban soils to the atmosphere. *Ecol. Appl.* 14:975-981.
- Lewis, J.D. and D.J. Bremer. 2013. Different nitrogen management regimes affect nitrous oxide emissions among one cool-season and two warm-season turfgrasses. *Int. Turf. Soc. Res. J.* 12:31-38.



Figure 1. Automated rainout shelter moving across plots activated by 0.01 inch of rain.

K-STATE
Research and Extension

Kansas State University
Agricultural Experiment Station
and Cooperative Extension Service





Figure 2. Plots received precise irrigation amounts based on daily ET during summer period.



Figure 3. Close-up of one of twelve static chambers used for sampling N_2O .






Figure 4. Cumulative fluxes of N₂O-N from plots treated with urea, polymer-coated urea, and untreated. Vertical dashed lines represent fertilization dates.




Figure 5. Visual quality ratings (9 = best quality) of ‘Meyer’ zoysiagrass prior to and following the summer period under the automated rainout shelter. Plots were fertilized with urea and poly on DOY 153 (2 June) and only urea fertilizer again on DOY 197 (July 16). Any points below yellow horizontal dashed line at rating of 6 would be below acceptable quality. Means followed by the same letter on a date are not significantly different according to Fisher’s protected least significant difference test ($P \leq 0.05$).

